

HEALTHWEST

Practice Guideline

No. 12-010

Prepared by:
Pharmacy Work Group

Effective: February 1, 2006
Revised: April 10, 2024

Approved by:

Subject: Simultaneous Use of
Multiple Psychotropic Medications

DocuSigned by:

Rich Francisco

Rich Francisco

Executive Director

I. PURPOSE

To encourage the use of monotherapy with psychotropic agents in a pharmaco-therapeutic regimen.

II. APPLICATION

HealthWest Physicians, Physician Assistants, and Nurse Practitioners.

III. DEFINITIONS

Psychotropic Medications: Medications listed in the Physician's Desk Reference (PDR) that effect the functioning of brain chemistry.

Simultaneous Use of Multiple Medications: Two (2) or more medications from the same class being prescribed at the same time. (Exception: Use of a depot anti-psychotic agent plus a different oral agent as a PRN [as needed] may in some cases be justified.)

IV. PROTOCOL

A. Continuous Review of Medications by Prescribers

1. Current medications will be reviewed by the prescriber for individuals seen in outpatient clinics.
2. The prescriber will implement a specific plan to eliminate excess psychotropic agents in cases of prescription of multiple psychotropic medications.

B. Documentation

1. The prescriber will document in each case where simultaneous use of multiple psychotropic medications are in use.
 - a. Documentation will be in the Treatment Plan Section of the Psychiatric Evaluation or Medication Review.

- b. Documentation will specify *either*:
 - (i) The *rationale* for continuing multiple psychotropic use, or
 - (ii) The *specific plan* for eliminating superfluous psychotropic agents through gradual dosage titrations. Reference to published algorithms for psychotropic therapy is suggested.
2. Random peer reviews will occur in cases of simultaneous use of multiple psychotropic use when two (2) or more medications are prescribed from the same class.

BD/hb